

# Air quality bulletin

---

North Queensland

November 2015

## Prepared by

Air Quality Monitoring  
Environmental Monitoring and Assessment Sciences  
Science Division  
Department of Science, Information Technology and Innovation  
PO Box 5078  
Brisbane QLD 4001

© The State of Queensland (Department of Science, Information Technology and Innovation) 2016

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 3.0 Australia (CC BY) licence



Under this licence you are free, without having to seek permission from DSITI, to use this publication in accordance with the licence terms.

You must keep intact the copyright notice and attribute the State of Queensland, Department of Science, Information Technology and Innovation as the source of the publication.

For more information on this licence visit <http://creativecommons.org/licenses/by/3.0/au/deed.en>

## Disclaimer

This document has been prepared with all due diligence and care, based on the best available information at the time of publication. The department holds no responsibility for any errors or omissions within this document. Any decisions made by other parties based on this document are solely the responsibility of those parties. Information contained in this document is from a number of sources and, as such, does not necessarily represent government or departmental policy.

If you need to access this document in a language other than English, please call the Translating and Interpreting Service (TIS National) on 131 450 and ask them to telephone Library Services on +61 7 3170 5725.

## Acknowledgements

This report has been prepared by the Department of Science, Information Technology and Innovation. Acknowledgement is made of the following organisations that have supplied air quality monitoring data for this bulletin:

- Sun Metals Corporation
- Port of Townsville Limited

January 2016

## Introduction

Air quality monitoring gathers information on the quality of the air environment. The objectives of the monitoring are to check compliance with ambient air quality guidelines, identify long-term trends in air quality, investigate local air quality concerns, and assess the effectiveness of air quality management strategies.

Air quality monitoring was carried out by the Queensland Government at three sites in Townsville and two sites in Mount Isa during November 2015. In addition, monitoring was also conducted by the Port of Townsville Limited and Sun Metals Corporation in Townsville. In May 2014 the former Townsville Port monitoring site operated by Port of Townsville Limited and the Queensland Government's Townsville Coast Guard monitoring site were amalgamated into a single monitoring site located at the Townsville Coast Guard.

Air pollutants monitored included ozone, nitrogen dioxide, sulfur dioxide, PM<sub>10</sub> (particles less than 10µm in diameter), TSP (total suspended particulate matter - particles less than 50µm approximately in diameter) and dustfall large enough to settle from the air. The TSP and dustfall samples were analysed for metal content.

The air pollutants monitored at North Queensland sites are shown in Table 1. Site locations are shown in Figure 23 on the last page of this bulletin.

Table 1. Air pollutants monitored at North Queensland sites

	Ozone	Nitrogen dioxide	Sulfur dioxide	PM <sub>10</sub>	TSP	Lead	Copper	Zinc	Nickel	Arsenic	Cadmium	Dustfall	Dustfall lead
<b>Townsville</b>													
Pimlico	✓	✓	✓	✓								✓	✓
Coast Guard				✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
North Ward (Warburton St)												✓	✓
Stuart (industry)			✓										
<b>Mount Isa</b>													
Menzies			✓										
The Gap			✓	✓		✓				✓	✓		

## Reporting protocol

Data presented in this bulletin are based on clock hours. Hourly or other averages are constrained to start and finish on a clock hour.

## Air quality summary graphs

Figures 1 to 14 summarise available air quality data for sampling days at Townsville sites during November 2015. Monthly average dustfall and deposited lead for Townsville sites are shown in figures 15 and 16. Figures 17 to 20 summarise air quality data for sampling days at Mount Isa sites during November. The maximum recorded level for each day is used to show the day-to-day variation in air quality.

## Air quality summary tables

Tables 5 to 22 present monthly summaries of air quality data for the preceding 12 months. These tables show the month-to-month variation in air quality. A monthly entry is given when at least three-fifths of the maximum possible number of observations during the month are available. When data is not available for the entire month, due to equipment malfunction, this is indicated by the abbreviation 'n.d.' (no data). A dash is inserted when less than three-fifths of data are available. Where no data is recorded, the reason for the low data availability is summarised in Table 23 at the end of this bulletin.

## Guidelines

Wherever possible, air quality measurements are compared against Australian air quality standards. In their absence relevant international standards are used for comparison.

Measured concentrations of nitrogen dioxide, ozone, sulfur dioxide, PM<sub>10</sub>, TSP, lead, nickel, cadmium and arsenic are compared to the air quality objectives contained in the Queensland *Environmental Protection (Air) Policy 2008* (EPP (Air)) to assess whether pollutant levels could harm health and well being.

Sulfur dioxide, PM<sub>10</sub>, arsenic and cadmium levels in Mount Isa are also compared against the air quality limits specified in the *Transitional Environmental Program (TEP) Xstrata Mount Isa Mines (Air Quality)*.

Copper and zinc levels present in TSP particles are compared against the Ontario Ministry of Environment ambient air quality criteria.

Limit values for TSP and dustfall specified in the Department of Environment and Heritage Protection (DEHP) guideline document *Application requirements for activities with impacts to Air* (Air Impacts Guideline) are used to assess dust nuisance levels.

The relevant guidelines are shown in the air quality summary table for each pollutant.

## Compliance with air quality guidelines - Townsville

During November, measured pollutant levels, with the exception of dustfall, did not exceed the relevant air quality guideline at Queensland Government and industry air monitoring sites in Townsville.

The DEHP dust nuisance limit value for dustfall was exceeded at the North Ward monitoring site during November. This exceedence follows an extended period of low rainfall which led to local activities (such as grounds maintenance at the Sports Reserve) producing greater amounts of dust in the vicinity of the sampler. Lower dust levels at the closer Townsville Coast Guard monitoring site during the same period indicates that Port of Townsville activities were not responsible for the high dustfall levels at the North Ward site.

Table 2. Number of occasions during November when measured levels exceeded EPP (Air) air quality objectives for nitrogen dioxide, ozone, sulfur dioxide, PM<sub>10</sub>, TSP, lead, nickel, arsenic and cadmium; Ontario Ministry of Environment air quality criteria for copper and zinc; and DEHP nuisance dust limits for TSP and dustfall at Queensland Government and industry air monitoring sites in Townsville.

Pollutant	Averaging Period	Exceedences
Nitrogen dioxide	<i>EPP(Air)</i> Annual	0
	1-hour	0
Ozone	<i>EPP(Air)</i> 4-hour	0
	1-hour	0
Sulfur dioxide	<i>EPP(Air)</i> Annual	0
	24-hour	0
	1-hour	0
PM <sub>10</sub>	<i>EPP(Air)</i> 24-hour	0
	TSP (24-hour period refers to dust nuisance, not health and well being)	<i>EPP(Air)</i> Annual
<i>DEHP limit</i> 24-hour		0
TSP Lead	<i>EPP(Air)</i> Annual	0
	TSP Copper	<i>Ontario criterion</i> 24-hour
TSP Zinc	<i>Ontario criterion</i> 24-hour	0
PM <sub>10</sub> Nickel	<i>EPP(Air)</i> Annual	0
PM <sub>10</sub> Arsenic	<i>EPP(Air)</i> Annual	0
	PM <sub>10</sub> Cadmium	<i>EPP(Air)</i> Annual
Dustfall (30-day period refers to dust nuisance, not health and wellbeing)	<i>DEHP limit</i> 30-day	1

## Compliance with air quality guidelines - Mount Isa

During November, measured pollutant levels, with the exception of sulfur dioxide, PM<sub>10</sub> and arsenic, did not exceed the relevant air quality guideline at Queensland Government air monitoring sites in Mount Isa.

From April 2012 smelter operations in Mount Isa have been operating under a Transitional Environmental Program (TEP) which sets out a staged program of works to achieve compliance with air quality objectives set out in the *Environmental Protection (Air) Policy 2008* by December 2016. Currently the TEP requires smelter operations to achieve compliance with the limits in Table 4 at industry air monitoring sites in the Mount Isa community (note: the Queensland Government monitoring sites are not part of this network).

Sulfur dioxide levels exceeded the EPP (Air) 1-hour objective for a total of seven 1-hour periods at the Menzies monitoring site and six 1-hour periods at The Gap monitoring sites during November due to industry emissions. Sulfur dioxide levels at both the Menzies and The Gap monitoring sites complied with the TEP limits during November.

There have been no exceedences of the TEP 1-hour sulfur dioxide limit value at either the Menzies or The Gap monitoring sites since 1 January 2015.

Windblown dust generated by the passage of a low pressure trough led to an exceedence of the EPP (Air) objective / TEP limit value for PM<sub>10</sub> at The Gap monitoring site on 27 November. On this day winds blew predominantly from the south-east, indicating

Table 3. Number of occasions during November when measured levels exceeded EPP (Air) air quality objectives for sulfur dioxide, PM<sub>10</sub>, lead, arsenic and cadmium; and the TEP limits for sulfur dioxide, PM<sub>10</sub>, arsenic and cadmium at Queensland Government air monitoring sites in Mount Isa.

Pollutant	Averaging Period	Exceedences
Sulfur dioxide	<i>EPP (Air)</i>	
	Annual	0
	24-hour	0
	1-hour	13
	<i>TEP</i>	
PM <sub>10</sub>	Annual	0
	1-hour	0
PM <sub>10</sub>	<i>EPP (Air) / TEP</i>	
	24-hour	1
TSP Lead	<i>EPP (Air)</i>	
	Annual	0
PM <sub>10</sub> Arsenic	<i>EPP (Air)</i>	
	Annual	1
	<i>TEP</i>	
PM <sub>10</sub> Cadmium	Annual	0
	<i>TEP</i>	
	Annual	0

Table 4. Transitional Environmental Program (TEP) air quality limits currently applying to smelter operations in Mount Isa

Pollutant	Averaging period	Limit value	Number of allowable exceedences	Period when limit value applies
Sulfur dioxide	Annual	57 µg/m <sup>3</sup> <sup>(a)</sup> (= 0.020 ppm)	Nil	21 April 2012 to 31 December 2016
	1-hour	2500 µg/m <sup>3</sup> <sup>(b)</sup> (= 0.955 ppm)	3 events per calendar year	
PM <sub>10</sub>	24-hour	50 µg/m <sup>3</sup>	7 days	1 January to 31 December 2015
Arsenic	Annual	0.017 µg/m <sup>3</sup>	Nil	1 January to 31 December 2015
Cadmium	Annual	0.007 µg/m <sup>3</sup>	Nil	1 January 2014 to 31 December 2016

<sup>(a)</sup> measured at 0°C, <sup>(b)</sup> measured at 25°C

that industry emissions were unlikely to have contributed significantly to this exceedence. There have been five exceedences of the TEP PM<sub>10</sub> limit value at The Gap site since 1 January 2015. Only two of these exceedences have occurred during wind conditions where industry emissions may have contributed to the exceedence.

Lead levels at The Gap monitoring site over the 12-month period ending November 2015 complied with the EPP (Air) objective.

The average PM<sub>10</sub> arsenic concentration at The Gap monitoring site over the 12-month period ending November 2015 exceeded the EPP (Air) objective but complied with the TEP limit value.

Cadmium levels at The Gap monitoring site over the 12-month period ending November 2015 complied with both the EPP (Air) objective and the TEP limit value.

## Measured ambient concentrations - Townsville

### Nitrogen dioxide

Figure 1. Ambient concentrations of nitrogen dioxide at the Pimlico site. Daily maximum 1-hour average concentrations (ppm), November 2015.




Table 5. Ambient concentrations of nitrogen dioxide. Annual average and monthly maximum 1-hour concentrations (ppm), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Pimlico</b>												
Annual average:	0.004											
Maximum 1-hour	0.017	0.012	0.014	0.014	0.031	0.028	0.028	0.031	0.040	0.025	0.015	0.011
% I.A.	98	79	98	98	98	94	98	98	97	98	98	98
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.												
The <i>Environmental Protection (Air) Policy 2008</i> air quality objectives for nitrogen dioxide are an annual average of 0.030 ppm and a 1-hour average of 0.120 ppm (not to be exceeded on more than one day per year).												

### Ozone (photochemical oxidants)

Figure 2. Ambient concentrations of ozone at the Pimlico site. Daily maximum 4-hour average concentrations (ppm), November 2015.




Figure 3. Ambient concentrations of ozone at the Pimlico site. Daily maximum 1-hour average concentrations (ppm), November 2015.




Table 6. Ambient concentrations of ozone. Monthly maximum 4-hour and 1-hour concentrations (ppm), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Pimlico</b>												
Maximum 4-hour	0.044	0.035	0.031	0.033	0.037	0.036	0.042	0.041	0.043	0.049	0.039	0.035
Maximum 1-hour	0.045	0.037	0.033	0.034	0.040	0.040	0.045	0.043	0.049	0.051	0.043	0.040
% I.A.	98	98	98	98	98	94	98	98	97	98	96	98
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available. The <i>Environmental Protection (Air) Policy 2008</i> air quality objectives for ozone are a 4-hour average of 0.080 ppm (not to be exceeded on more than one day per year) and a 1-hour average of 0.100 ppm (not to be exceeded on more than one day per year).												

**Sulfur dioxide**

Figure 4. Ambient concentrations of sulfur dioxide at Pimlico and Stuart sites. Daily 24-hour average concentrations (ppm), November 2015.




Figure 5. Ambient concentrations of sulfur dioxide at Pimlico and Stuart sites. Daily maximum 1-hour average concentrations (ppm), November 2015.




Table 7. Ambient concentrations of sulfur dioxide. Annual average and monthly maximum 24-hour and 1-hour average concentrations (ppm), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Pimlico</b>												
Annual average:	0.001											
Maximum 24-hour	0.002	0.001	0.003	0.002	0.002	0.003	0.001	0.001	0.001	0.001	0.001	0.000
Maximum 1-hour	0.004	0.003	0.004	0.003	0.004	0.004	0.003	0.004	0.003	0.003	0.002	0.001
% I.A.	98	98	98	98	98	94	98	98	97	98	98	98
<b>Stuart (industry-operated site)</b>												
Annual average:	-											
Maximum 24-hour	0.001	-	0.001	0.001	0.001	-	n.d.	n.d.	0.001	0.002	0.000	0.002
Maximum 1-hour	0.002	-	0.002	0.003	0.002	-	n.d.	n.d.	0.005	0.004	0.003	0.003
% I.A.	96	40	64	80	91	51	0	0	87	98	98	92
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.												
The <i>Environmental Protection (Air) Policy 2008</i> air quality objectives for sulfur dioxide are an annual average of 0.020 ppm, a 24-hour average of 0.080 ppm (not to be exceeded on more than one day per year) and a 1-hour average of 0.200 ppm (not to be exceeded on more than one day per year).												



**PM<sub>10</sub>**

Figure 6. Ambient concentrations of PM<sub>10</sub> at Pimlico and Coast Guard sites. Daily 24-hour average concentrations (µg/m<sup>3</sup>), November 2015.




Table 8. Ambient concentrations of PM<sub>10</sub>. Monthly maximum 24-hour average concentrations (µg/m<sup>3</sup>), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Pimlico</b>												
Maximum 24-hour	21.9	20.7	25.5	21.1	21.9	25.7	30.0	28.8	31.8	36.6	42.0	25.7
% I.A.	99	95	92	87	88	94	99	100	98	97	86	95
<b>Coast Guard</b>												
Maximum 24-hour	28.8	30.6	28.7	29.7	37.5	38.1	34.9	31.6	30.5	35.7	35.0	26.5
% I.A.	90	89	96	98	94	97	100	99	100	88	98	99
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available. The <i>Environmental Protection (Air) Policy 2008</i> air quality objective for PM <sub>10</sub> is a 24-hour average of 50 µg/m <sup>3</sup> (not to be exceeded on more than five days per year).												

**TSP**

Figure 7. Ambient concentrations of TSP (continuous monitoring) at the Coast Guard site. Daily 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ), November 2015.




Figure 8. Ambient concentrations of TSP (one day in six monitoring) at the Coast Guard site. 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ), November 2015.




Table 9. Ambient concentrations of TSP. Monthly maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard (continuous monitoring)</b>												
Annual average:	28.4											
Maximum 24-hour	39.8	44.0	45.8	43.9	60.4	71.9	41.4	56.6	53.8	55.8	73.9	34.6
% I.A.	100	89	97	98	94	96	90	93	99	89	94	99
<b>Coast Guard (one day in six monitoring)</b>												
Annual average:	36.1											
Maximum 24-hour	59.3	40.2	44.4	55.2	63.3	57.0	44.9	48.8	64.5	76.7	64.4	33.1
% I.A.	100	100	100	100	100	100	80	100	80	67	100	100

% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.

The *Environmental Protection (Air) Policy 2008* air quality objective for TSP is an annual average of  $90 \mu\text{g}/\text{m}^3$ .

The Department of Environment and Heritage Protection Air Impacts Guideline recommends that short-term (24-hour) TSP concentrations be compared against the trigger levels provided in the New Zealand Ministry for the Environment's *Good practice guide for assessing and managing the environmental effects of dust emissions to assess dust nuisance impacts*. The New Zealand dust nuisance (sensitive areas) trigger level is a 24-hour average of  $80 \mu\text{g}/\text{m}^3$ .

**TSP Lead**

Figure 9. Ambient concentrations of TSP lead (one day in six monitoring) at the Coast Guard site. Annual average concentration ( $\mu\text{g}/\text{m}^3$ ), December 2014 to November 2015.




Table 10. Ambient concentrations of TSP lead. Annual average and monthly maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ) for one day in six monitoring, December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Annual average:	0.14											
Maximum 24-hour	0.79	0.21	0.17	0.62	0.43	0.11	0.21	0.11	0.10	0.24	0.19	1.29
% I.A.	100	100	100	100	100	100	80	100	80	67	100	100
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.												
The <i>Environmental Protection (Air) Policy 2008</i> air quality objective for lead is an annual average of $0.5 \mu\text{g}/\text{m}^3$ .												
The limit of reporting is the minimum measured lead concentration that can be determined with the sampling equipment and laboratory method used. Lead concentrations below this limit are preceded by a "<" sign in the table. The annual average concentration has been calculated using half the minimum measurable concentration value for samples where no lead was detected.												

### TSP Copper

Figure 10. Ambient concentrations of TSP copper (one day in six monitoring) at the Coast Guard site. Maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ), November 2015.




Table 11. Ambient concentrations of TSP copper. Monthly maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ) for one day in six monitoring, December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Maximum 24-hour	0.02	0.02	0.08	1.71	0.91	0.05	0.04	0.07	0.18	0.07	0.04	0.02
% I.A.	100	100	100	100	100	100	80	100	80	67	100	100
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available. The Ontario Ministry of Environment ambient air quality criterion for copper is a 24-hour average of $50 \mu\text{g}/\text{m}^3$ . The limit of reporting is the minimum measured copper concentration that can be determined with the sampling equipment and laboratory method used. Copper concentrations below this limit are preceded by a "<" sign in the table.												

**TSP Zinc**

Figure 11. Ambient concentrations of TSP zinc (one day in six monitoring) at the Coast Guard site. Maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ), November 2015.




Table 12. Ambient concentrations of TSP zinc. Monthly maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ) for one day in six monitoring, December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Maximum 24-hour	1.12	0.46	1.90	3.16	2.28	1.60	1.90	0.57	0.55	3.09	0.86	0.64
% I.A.	100	100	100	100	100	100	80	100	80	67	100	100
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available. The Ontario Ministry of Environment ambient air quality criterion for zinc is a 24-hour average of $120 \mu\text{g}/\text{m}^3$ . The limit of reporting is the minimum measured zinc concentration that can be determined with the sampling equipment and laboratory method used. Zinc concentrations below this limit are preceded by a "<" sign in the table.												

**TSP Nickel**

Figure 12. Ambient concentrations of TSP nickel (one day in six monitoring) at the Coast Guard site. Annual average concentrations ( $\mu\text{g}/\text{m}^3$ ), December 2014 to November 2015.




Table 13. Ambient concentrations of TSP nickel. Annual average and monthly maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ) for one day in six monitoring, December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Annual average:	0.015											
Maximum 24-hour	0.098	0.033	0.040	0.052	0.027	<0.010	0.016	<0.010	0.054	0.033	0.047	0.047
% I.A.	100	100	100	100	100	100	80	100	80	67	100	100

% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.

The *Environmental Protection (Air) Policy 2008* air quality objective for nickel is an annual average of  $0.020 \mu\text{g}/\text{m}^3$  (measured as the total metal content in  $\text{PM}_{10}$  particles).

Monitoring conducted by the Queensland Government measures the amount of nickel present in the TSP fraction. As  $\text{PM}_{10}$  is a subset of TSP, if the TSP nickel concentration is less than the EPP (Air)  $\text{PM}_{10}$  objective value, it follows that the  $\text{PM}_{10}$  nickel concentration complies with the EPP (Air) objective.

The limit of reporting is the minimum measured nickel concentration that can be determined with the sampling equipment and laboratory method used. Nickel concentrations below this limit are preceded by a "<" sign in the table. Annual average concentrations have been calculated using half the minimum measurable concentration value for samples where no nickel was detected.

**TSP Arsenic**

Figure 13. Ambient concentrations of TSP arsenic (one day in six monitoring) at the Coast Guard site. Annual average concentrations ( $\mu\text{g}/\text{m}^3$ ), December 2014 to November 2015.




Table 14. Ambient concentrations of TSP arsenic. Monthly maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ) for one day in six monitoring, December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Annual average	0.001											
Maximum 24-hour	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
% I.A.	100	100	100	100	100	100	80	100	80	67	100	100

% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.

The *Environmental Protection (Air) Policy 2008* air quality objective for arsenic is an annual average of  $0.006 \mu\text{g}/\text{m}^3$  (measured as the total metal content in  $\text{PM}_{10}$  particles).

Monitoring conducted by the Queensland Government measures the amount of arsenic present in the TSP fraction. As  $\text{PM}_{10}$  is a subset of TSP, if the TSP arsenic concentration is less than the EPP (Air)  $\text{PM}_{10}$  objective value, it follows that the  $\text{PM}_{10}$  arsenic concentration complies with the EPP (Air) objective.

The limit of reporting is the minimum measured arsenic concentration that can be determined with the sampling equipment and laboratory method used. Arsenic concentrations below this limit are preceded by a "<" sign in the table. Annual average concentrations have been calculated using half the minimum measurable concentration value for samples where no arsenic was detected.

### TSP Cadmium

Figure 14. Ambient concentrations of TSP cadmium (one day in six monitoring) at the Coast Guard site. Annual average concentrations ( $\mu\text{g}/\text{m}^3$ ), December 2014 to November 2015.




Table 15. Ambient concentrations of TSP cadmium. Monthly maximum 24-hour average concentrations ( $\mu\text{g}/\text{m}^3$ ) for one day in six monitoring, December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Annual Average	0.0020											
Maximum 24-hour	0.005	0.002	0.005	0.015	0.007	0.004	0.006	0.002	0.002	0.009	0.003	0.004
% I.A.	100	100	100	100	100	100	80	100	80	67	100	100

% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.

The *Environmental Protection (Air) Policy 2008* air quality objective for cadmium is an annual average of  $0.005 \mu\text{g}/\text{m}^3$  (measured as the total metal content in  $\text{PM}_{10}$  particles).

Monitoring conducted by the Queensland Government measures the amount of cadmium present in the TSP fraction. As  $\text{PM}_{10}$  is a subset of TSP, if the TSP cadmium concentration is less than the EPP (Air)  $\text{PM}_{10}$  objective value, it follows that the  $\text{PM}_{10}$  cadmium concentration complies with the EPP (Air) objective.

The limit of reporting is the minimum measured cadmium concentration that can be determined with the sampling equipment and laboratory method used. Cadmium concentrations below this limit are preceded by a "<" sign in the table. Annual average concentrations have been calculated using half the minimum measurable concentration value for samples where no cadmium was detected.



## Dustfall

Figure 15. Dustfall monitoring at the Coast Guard, North Ward (Warburton Street) and Pimlico sites. Daily average dust (insoluble solids fraction) deposition rate ( $\text{mg}/\text{m}^2/\text{day}$ ) for month of November 2015.




Table 16. Daily average dust (insoluble solids fraction) deposition rate ( $\text{mg}/\text{m}^2/\text{day}$ ), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Daily average	n.d.	31	56	45	47	53	46	30	46	241	80	91
<b>North Ward (Warburton St)</b>												
Daily average	111	17	111	123	141	217	171	158	204	197	n.d.	165
<b>Pimlico</b>												
Daily average	22	n.d.	30	10	13	20	11	7	40	21	17	15
n.d. indicates no data are available.												
The Department of Environment and Heritage Protection Air Impacts Guideline recommends a dust deposition limit of $120 \text{ mg}/\text{m}^2/\text{day}$ , averaged over one month, be used to assess dust nuisance.												
The limit of reporting is the minimum measured dust deposition rate that can be determined with the sampling equipment and laboratory method used. Dust deposition rates below this limit are preceded by a "<" sign in the table.												

**Dustfall lead**

Figure 16. Dustfall lead monitoring at the Coast Guard, North Ward (Warburton Street) and Pimlico sites. Daily average lead deposition rate ( $\mu\text{g}/\text{m}^2/\text{day}$ ) for month of November 2015.




Table 17. Daily average lead deposition rate ( $\mu\text{g}/\text{m}^2/\text{day}$ ), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Townsville</b>												
<b>Coast Guard</b>												
Annual average:	305											
Daily average	n.d.	314	481	387	313	167	136	85	154	262	606	400
<b>North Ward (Warburton St)</b>												
Annual average:	<30											
Daily average	<37	<29	<37	<32	<31	<33	<36	<30	<36	<34	n.d.	<29
<b>Pimlico</b>												
Annual average:	<30											
Daily average	<37	n.d.	<37	<32	<31	<33	<36	<24	<50	<34	<29	<29

n.d. indicates no data are available.

There is no air quality objective for ambient lead deposition. Some data indicate that lead fallout levels between 250 and 750  $\mu\text{g}/\text{m}^2/\text{day}$  (averaged over a 12-month period) are associated with a slight increase in blood lead levels (Air Quality Guidelines for Europe, Second Edition, World Health Organization, 2000).

The limit of reporting is the minimum measured lead deposition rate that can be determined with the sampling equipment and laboratory method used. Lead deposition rates below this limit are preceded by a "<" sign in the table.

## Measured ambient concentrations - Mount Isa

### Sulfur dioxide

Figure 17. Ambient concentrations of sulfur dioxide at the Menzies and The Gap sites. Daily 24-hour average concentrations (ppm), November 2015.




Figure 18. Ambient concentrations of sulfur dioxide at the Menzies and The Gap sites. Daily maximum 1-hour average concentrations (ppm), November 2015.




Table 18. Ambient concentrations of sulfur dioxide. Annual average and monthly maximum 24-hour and 1-hour average concentrations (ppm), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Mount Isa</b>												
<b>Menzies</b>												
Annual average:	0.007											
Maximum 24-hour	0.096	0.081	0.106	0.045	0.037	0.031	0.006	0.047	0.035	0.037	0.040	0.079
Maximum 1-hour	0.497	0.512	0.428	0.468	0.276	0.245	0.050	0.577	0.285	0.216	0.371	0.466
% I.A.	97	98	98	98	98	98	98	98	98	98	97	98
<b>The Gap</b>												
Annual average:	0.005											
Maximum 24-hour	0.059	0.045	0.035	0.036	0.016	0.034	0.004	0.030	0.022	0.021	0.056	0.044
Maximum 1-hour	0.568	0.341	0.231	0.383	0.123	0.340	0.067	0.335	0.212	0.141	0.463	0.494
% I.A.	97	98	98	98	98	98	98	98	98	98	96	98
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available. The <i>Environmental Protection (Air) Policy 2008</i> air quality objectives for sulfur dioxide are an annual average of 0.020 ppm, a 24-hour average of 0.080 ppm (not to be exceeded on more than one day per year) and a 1-hour average of 0.200 ppm (not to be exceeded on more than one day per year). The Transitional Environmental Program (TEP) air quality limits for sulfur dioxide are an annual average of 0.020 ppm and a maximum 1-hour average of 0.955 ppm, not to be exceeded more than three times during 2015.												

**PM<sub>10</sub>**

Figure 19. Ambient concentrations of PM<sub>10</sub> at The Gap site. Daily 24-hour average concentrations (µg/m<sup>3</sup>), November 2015.




Table 19. Ambient concentrations of PM<sub>10</sub>. Monthly maximum 24-hour average concentrations (µg/m<sup>3</sup>), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Mount Isa</b>												
<b>The Gap</b>												
Maximum 24-hour	91.5	65.3	45.7	47.7	37.7	56.9	29.0	37.9	50.0	59.7	40.6	54.2
% I.A.	98	96	97	99	100	100	93	100	99	100	99	100
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available. The <i>Environmental Protection (Air) Policy 2008</i> air quality objective for PM <sub>10</sub> is a 24-hour average of 50 µg/m <sup>3</sup> (not to be exceeded on more than five days per year). The Transitional Environmental Program air quality limit for PM <sub>10</sub> is a maximum 24-hour average of 50 µg/m <sup>3</sup> , not to be exceeded on more than seven days between 1 January 2015 and 31 December 2015.												

### TSP Lead

Figure 20. Ambient concentrations of TSP lead at The Gap site. Annual average concentration (µg/m<sup>3</sup>), December 2014 to November 2015.




Table 20. Ambient concentrations of TSP lead. Annual average and monthly maximum 24-hour average concentrations (µg/m<sup>3</sup>), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Mount Isa</b>												
<b>The Gap</b>												
Annual average:	0.10											
Maximum 24-hour	0.69	0.27	0.39	0.23	0.02	0.02	0.03	0.03	0.14	0.10	0.15	1.28
% I.A.	100	100	100	100	100	100	100	100	100	100	100	100
% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available. The <i>Environmental Protection (Air) Policy 2008</i> air quality objective for lead is an annual average of 0.5 µg/m <sup>3</sup> . The limit of reporting is the minimum measured lead concentration that can be determined with the sampling equipment and laboratory method used. Lead concentrations below this limit are preceded by a "<" sign in the table. The annual average concentration has been calculated using half the minimum measurable concentration value for samples where no lead was detected.												

**PM<sub>10</sub> Arsenic**

Figure 21. Ambient concentrations of PM<sub>10</sub> arsenic at The Gap site. Annual average concentration (µg/m<sup>3</sup>), December 2014 to November 2015.




Table 21. Ambient concentrations of PM<sub>10</sub> arsenic. Annual average and monthly maximum 24-hour average concentrations (µg/m<sup>3</sup>), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Mount Isa</b>												
<b>The Gap</b>												
Annual average:	0.007											
Maximum 24-hour	0.066	0.039	0.022	0.022	<0.001	<0.001	<0.001	<0.001	0.019	0.008	0.028	0.046
% I.A.	100	100	100	100	100	100	100	100	100	100	100	100

% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.

The *Environmental Protection (Air) Policy 2008* air quality objective for arsenic is an annual average of 0.006 µg/m<sup>3</sup> (measured as the total metal content in PM<sub>10</sub> particles).

The Transitional Environmental Program air quality limit for arsenic is an annual average of 0.017 µg/m<sup>3</sup> (measured as the total metal content in PM<sub>10</sub> particles between 1 January and 31 December 2015).

Monitoring conducted by the Queensland Government measures the amount of arsenic present in the TSP fraction (the TSP fraction is collected so lead levels can be compared against the EPP (Air) objective). Monitoring using co-located TSP and PM<sub>10</sub> high volume samplers has determined that the ratio of PM<sub>10</sub> arsenic to TSP arsenic is 0.88:1 in Mount Isa. The PM<sub>10</sub> arsenic values presented in this table have been generated by applying this factor to the measured TSP arsenic concentrations.

The limit of reporting is the minimum measured arsenic concentration that can be determined with the monitoring instrumentation used. Arsenic concentrations below this limit are preceded by a "<" sign in the table. The annual average concentration has been calculated using half the minimum measurable concentration value for samples where no arsenic was detected.

**PM<sub>10</sub> Cadmium**

Figure 22. Ambient concentrations of PM<sub>10</sub> cadmium at The Gap site. Annual average concentration (µg/m<sup>3</sup>), December 2014 to November 2015.




Table 22. Ambient concentrations of PM<sub>10</sub> cadmium. Annual average and monthly maximum 24-hour average concentrations (µg/m<sup>3</sup>), December 2014 to November 2015.

Site	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<b>Mount Isa</b>												
<b>The Gap</b>												
Annual average:	0.001											
Maximum 24-hour	0.012	0.004	0.007	0.004	<0.001	<0.001	0.001	<0.001	0.002	<0.001	0.003	0.010
% I.A.	100	100	100	100	100	100	100	100	100	100	100	100

% I.A. indicates instrument availability. - indicates less than three-fifths of the data are available. n.d. indicates no data are available.

The *Environmental Protection (Air) Policy 2008* air quality objective for cadmium is an annual average of 0.005 µg/m<sup>3</sup> (measured as the total metal content in PM<sub>10</sub> particles).

The Transitional Environmental Program air quality limit for cadmium is an annual average of 0.007 µg/m<sup>3</sup> (measured as the total metal content in PM<sub>10</sub> particles between 1 January and 31 December 2015).

Monitoring conducted by the Queensland Government measures the amount of cadmium present in the TSP fraction (the TSP fraction is collected so lead levels can be compared against the EPP(Air) objective). Monitoring using co-located TSP and PM<sub>10</sub> high volume samplers has determined that the ratio of PM<sub>10</sub> cadmium to TSP cadmium is 0.76:1 in Mount Isa. The PM<sub>10</sub> cadmium values presented in this table have been generated by applying this factor to the measured TSP cadmium concentrations.

The limit of reporting is the minimum measured cadmium concentration that can be determined with the monitoring instrumentation used. Cadmium concentrations below this limit are preceded by a "<" sign in the table. The annual average concentration has been calculated using half the minimum measurable concentration value for samples where no cadmium was detected.

**Data Availability**

When required, Table 23 summarises the reasons for data availability below the minimum criteria for reporting at North Queensland monitoring sites.

Table 23. Reasons for low data availability at North Queensland ambient air monitoring sites during November 2015.

Station	Air Pollutant	Cause
Nil		

**Related air quality information**

Current hourly air quality data is available from the internet at [www.ehp.qld.gov.au/air/data/search.php](http://www.ehp.qld.gov.au/air/data/search.php).

Additional information on air quality monitoring and related issues is also available from the above website.

**Further information**

For further information about the data presented in this bulletin or related publications, contact:

Air Quality Monitoring  
 Environmental Monitoring and Assessment Sciences  
 Science Division  
 Department of Science, Information Technology and Innovation  
 Ecosciences Precinct  
 41 Boggo Rd  
 DUTTON PARK QLD 4102  
 Telephone (07) 3170 5477  
 Email: [air.sciences@dsitia.qld.gov.au](mailto:air.sciences@dsitia.qld.gov.au)

Figure 23. North Queensland ambient air quality monitoring station locations.

